TOBI® **Podhaler**® is an aminoglycoside antibiotic for Suppressive therapy of chronic pulmonary infection due to Pseudomonas aeruginosa in adults and children aged 6 years and older with **cystic fibrosis**¹







Tobramycin 28 mg inhalation powder hard capsules

Is the only inhaled dry powder tobramycin formulation that does not require a nebulizer in Israeli market (DPI)

THIS DRY POWDER FORMULATION PROVIDES:2

- Substantially improved intrapulmonary deposition efficiency
- Faster delivery
- · More convenient administration over nebulized formulations

SPECIAL PULMOSPHERE™ TECHNOLOGY^{2,3}

An emulsion-based spray drying process that produces spherical hollow-porous particles that delivers the antibiotic tobramycin in a dry powder form (**PulmoSphere™**) to the central and peripheral airways of the lungs

- Inhaled antibiotic therapy has proven efficacy to:4
- · Reduce pulmonary exacerbations
- Improve lung function
- Improve respiratory symptoms

If eradication fails and chronic infection with PA develops

DEMONSTRATED EFFICACY

- DPI tobramycin has better intrapulmonary deposition efficiency
 The deposition fraction for TIP (tobramycin inhalation powder) is ~ 3 times greater
 than with TIS (tobramycin inhalation solution)²
- Improvements in lung function, compared with placebo⁵
- Reduced P. aeruginosa sputum density⁵
- Reduced respiratory-related hospitalizations compared with placebo⁵
- Reduced use of additional antipseudomonal antibiotics compared with placebo⁵
- Delivers the high payload of drug to the lungs needed for effective treatment²
- The PulmoSphere™ particles are ideal for targeting the powder aerosol to the site of the PA infection in the airways²

DESIGNED FOR PATIENTS ON THE GO

- Increased Portability¹
- No refrigeration, room temperature store below 30°C
- · Store in the original package in order to protect from moisture
- · No power source required
- Ease of use (no nebulizer required)
- Quick Clean Up^{1,2}
- · No disinfection required wipe clean with a dry cloth

TOBI® PODHALER® ADMINISTRATION SCHEDULE¹

- Simple & rapid administration time of only 6 minutes:
- 4 capsules for inhalation the powder from all 4 capsules must be inhaled to receive the full dose of 112 mg
- · 2 times (AM & PM) as close to 12 hours apart, not less than 6 hours apart
- 6 minutes administration
- · 28 days on
- · 28 days off